

Anaphylaxis Emergency Action Plan

Patient Name:			Age:	
Allergies:				
Asthma ☐ Yes (high risk for sever		□No		
Additional health problems besides	anaphylaxis:			
Concurrent medications:				
MOUTH THROAT* SKIN GUT LUNG* HEART*	itching, swe itching, tigh itching, hiv vomiting, d shortness of	ms of Anaphylaxis elling of lips and/or tongo ntness/closure, hoarsene es, redness, swelling iarrhea, cramps of breath, cough, wheeze , dizziness, passing out	ss	
Only a few symptoms *Some s	may be prese ymptoms can	nt. Severity of symptoms be life-threatening. ACT	can change quickly. FAST!	
Emergency Action Steps - DO 1. Inject epinephrine in thigh using (ch	O NOT HESITA	ATE TO GIVE EPINEPHRI Adrenaclick (0.15 mg)	NE! Adrenaclick (0.3 mg)	
	[Auvi-Q (0.15 mg)	Auvi-Q (0.3 mg)	
	,	EpiPen Jr (0.15 mg)	☐ EpiPen (0.3 mg)	
		Epinephrine Injection, US (0.15 mg)	SP Auto-injector- authorized generi	
	[Other (0.15 mg)	Other (0.3 mg)	
Specify others:				
IMPORTANT: ASTHMA INHALERS A	ND/OR ANTIH	HISTAMINES CAN'T BE D	EPENDED ON IN ANAPHYLAXIS.	
2. Call 911 or rescue squad (before	calling contac	t)		
3. Emergency contact #1: home		work	cell	
Emergency contact #2: home		work	cell	
Emergency contact #3: home		work	cell	
Comments:				
,				
Doctor's Signature/Date/Phone Number	er			

This information is for general purposes and is not intended to replace the advice of a qualified health professional. For more information, visit www.aaaai.org. © 2017 American Academy of Allergy, Asthma & Immunology 4/2017

Parent's Signature (for individuals under age 18 yrs)/Date